Asthma care plan for education and care services

CONFIDENTIAL: Staff are trained in asthma first aid (see overleaf) and can provide routine asthma medication as authorised in this care plan by the treating doctor. Please advise staff in writing of any changes to this plan.

To be completed by the treating doctor and parent/quardian, for supervising staff and emergency medical personnel.

PLEASE PRINT CLEARLY

Photo of student (optional)

> Plan date / /201

Review date

			/201
Student's name	Date of birth		
Managing an asthma attack Staff are trained in asthma first aid (see	overleaf). Please write down anything different this student	might need if the	ey have an asthma attack:
Daily asthma management			
This student's usual asthma signs Cough Wheeze Difficulty breathing Other (please describe)	Frequency and severity Daily/most days Frequently (more than 5 x per year) Occasionally (less than 5 x per year) Other (please describe)		for this student's asthma (eg l'flu, smoke) — please detail:
Medication plan	thma medication? Yes N	lo lo lo lo acer/mask are su	pplied to staff.
Does this student need help to take ast Does this student use a mask with a sp *Does this student need a blue reliever Medication plan	thma medication? Yes Nacer? Puffer medication before exercise? Yes N	lo lo	pplied to staff. Time required
Does this student need help to take ast Does this student use a mask with a sp *Does this student need a blue reliever Medication plan If this student needs asthma medicatio	thma medication? Yes Nacer? Puffer medication before exercise? Yes Nacer? Nacer? Yes Nacer? Nacer. N	lo lo	
Does this student need help to take ast Does this student use a mask with a sp *Does this student need a blue reliever Medication plan If this student needs asthma medicatio	thma medication? Yes Nacer? Yes Note puffer medication before exercise? Yes Note puffer medication before exercise? Note puffer medication before exercise? Note puffer medication before exercise? Note puffer medication and specific puffer	lo lo lacer/mask are su	
Does this student need help to take ast Does this student use a mask with a sp *Does this student need a blue reliever Medication plan If this student needs asthma medicatio Name of medication and colour Doctor	thma medication? Yes Nacer? Yes Note puffer medication before exercise? Yes Note puffer medication before exercise? Note puffer medication before exercise? Note puffer medication before exercise? Note puffer medication and specific puffer medication before exercise? Parent/Guardian I have read, understood and agreed with this care plan and any attachments listed. I approve the release of this information to staff	lo lo lo lacer/mask are su Emergency co	Time required
Does this student need help to take ast Does this student use a mask with a sp *Does this student need a blue reliever Medication plan If this student needs asthma medicatio Name of medication and colour Doctor Name of doctor	thma medication? Yes Nacer? Yes Note puffer medication before exercise? Yes Note puffer medication before exercise? Yes Note puffer medication before exercise? Note puffer medication before exercise? Note puffer medication before exercise? Note puffer medication and specific puffer before the medication and specific puffer before puffs Note puffer medication before exercise? Note puffer medication and specific puffer before puffs Parent/Guardian I have read, understood and agreed with this care plan and any attachments listed. I approve the release of this information to staff and emergency medical personnel. I will notify the staff in writing if there are any changes to these instructions. I understand staff will seek emergency medical help as needed and that I am responsible	lo lo lo lo lacer/mask are su Emergency co Contact name	Time required









Asthma First Aid

1 Sit the person upright

- Be calm and reassuring
- Do not leave them alone



Give 4 separate puffs of blue/grey reliever puffer

- Shake puffer
- Put 1 puff into spacer
- Take <u>4 breaths</u> from spacer

Repeat until 4 puffs have been taken

Remember: Shake, 1 puff, 4 breaths

OR give 2 separate doses of a Bricanyl inhaler (age 6 & over) or a Symbicort inhaler (over 12)



Wait 4 minutes

If there is no improvement, give <u>4 more separate puffs of blue/grey reliever</u> as above

OR give 1 more dose of Bricanyl or Symbicort inhaler



If there is still no improvement call emergency assistance - Dial Triple Zero (000)

- Say 'ambulance' and that someone is having an asthma attack
- Keep giving <u>4 separate puffs</u> every <u>4 minutes</u> until emergency assistance arrives

OR give 1 dose of a Bricanyl or Symbicort every 4 minutes - up to 3 more doses of Symbicort



Call emergency assistance immediately - Dial Triple Zero (000)

- If the person is not breathing
- If the person's asthma suddenly becomes worse or is not improving
- If the person is having an asthma attack and a reliever is not available
- If you are not sure if it's asthma
- If the person is known to have Anaphylaxis follow their Anaphylaxis Action
 Plan, then give Asthma First Aid

Blue/grey reliever medication is unlikely to harm, even if the person does not have asthma



Call the **1800 ASTHMA** Helpline (1800 278 462) or visit asthmaaustralia.org.au

